



# Suspend<sup>®</sup> SC

## Insecticide

- Only for Use and Storage by Pest Control Operators and/or Commercial Applicators
- For Residential, Industrial and Institutional Control of Major Nuisance Pests Listed on The Label
- Controls Numerous Pests in and around Households and Other Structures
- Will not Damage Paints, Plastics, Fabrics or Other Surfaces Where Water Alone Causes No Damage
- For Insect Control on Commercial and Residential Turf and Landscape Ornamental Plants

**ACTIVE INGREDIENTS:**

\*Deltamethrin ..... 4.75%

**OTHER INGREDIENTS:** ..... 95.25%

100.00%

\*Contains 0.42 lb deltamethrin per gallon.

EPA Reg No. 432-763

### KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

Net Contents  
**1 Gallon**

4205919

4548174D

100504AV2

**BACKED**  
by **BAYER**<sup>™</sup>

FIRST AID	
If Inhaled	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-334-7577 for emergency medical treatment.	

FOR PRODUCT INFORMATION, CALL TOLL-FREE: 1-800-331-2867.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS & DOMESTIC ANIMALS

##### CAUTION

Harmful if inhaled. Avoid breathing vapor or spray mist. Contact with product may result in transient tingling and reddening of the skin. Remove contaminated clothing and wash before reuse.

##### ENVIRONMENTAL HAZARDS

This product is extremely toxic to fresh water and estuarine fish and invertebrates. Use with care when applying in areas adjacent to any body of water. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This pesticide is highly toxic to bees exposed to direct treatment. Do not apply or allow it to drift to crops or weeds on which bees are actively foraging.

##### PHYSICAL & CHEMICAL HAZARDS

Do not use water-based sprays of Suspend SC® Insecticide in conduits, motor housings, junction boxes, switch boxes, or other electrical equipment because of possible shock hazard.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

##### GENERAL INFORMATION:

Suspend SC Insecticide is a suspended concentrate insecticide containing 0.42 lb active ingredient per gallon. Suspend SC Insecticide is designed for use both indoors and outdoors and will provide effective knockdown and long residual control of the pests listed on this label.

##### APPLICATION RESTRICTIONS

Do not water the treated area to the point of run-off.

Do not make applications during rain.

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

- (1) Treatment to soil or vegetation around structures;
- (2) Applications to lawns, turf, and other vegetation;
- (3) Applications to building foundations, up to a maximum height of 3 feet.

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

**Mixing Instructions:** Suspend SC Insecticide is intended to be mixed with water and applied with hand pressurized or power operated sprayers, such as Actisol® or similar equipment, as a coarse spray or with conventional hydraulic boom-type sprayer or other suitable equipment for outdoor broadcast applications. Shake container well before using. When diluting, it is recommended to add approximate-

ly half the water to the spray tank, then add the proper amount of Suspend SC Insecticide in accordance with the rates in the appropriate Application Rate table. Agitate, then add the remaining water slowly to avoid excess foam. Thoroughly agitate the sprayer before using to insure proper suspension. If spraying is interrupted when using a back-pack or hand-held spray equipment, reinvert sprayer before use. If spray screens are used, they should be 50 mesh or larger. Dilute Suspend SC Insecticide only with water. Prepare spray solution when ready to make applications.

**Tank Mixing:** Suspend SC Insecticide may be tank-mixed with other currently registered pesticides unless expressly prohibited by the product label. A small volume mixing test with other products is recommended to ensure compatibility. Observe all restrictions and precautions, which appear on the labels of these products.

**Application Methods:** For indoor uses this product may be applied as crack and crevice, spot, general surface treatments. For residential building and structures apply only as a crack and crevice or spot treatment. Treat where pests are found or normally occur. For general pest control outdoors see this section for additional methods of application.

**Crack and Crevice Applications:** Use a low pressure system with a pinpoint or variable pattern nozzle to apply the spray mixture to areas such as floors, cracks and crevices in and around baseboards, walls, expansion joints, areas around water and sewer pipes, and voids formed by equipment or appliances.

**Spot Applications:** Use a low pressure system with a fan-type nozzle to uniformly apply spot treatments in and around cupboards, and between, beneath and behind equipment or appliances. For control of carpet beetles, fleas and ticks indoors, apply as a coarse, low pressure crack and crevice spot treatment to problem areas and infested habitats. Do not treat entire floor, carpet or floor coverings. Do not apply as a space spray. Application to furniture should be restricted to areas where prolonged contact by humans will not occur. Care should be taken to limit spot treatment to pet beds and infested floor coverings.

### Application Rates

Pests	Concentration of Active Ingredient <sup>1</sup>	Dilution Rate <sup>2</sup>
Ants, carpet beetles, firebrats, fleas, gnats, midges, centipedes, millipedes, pantry pests, silverfish, ticks(indoor)	0.01% - 0.03%w/w	0.25-0.75 fl oz/gal 12.5-37.5 fl oz/50 gals 25-75 fl oz/100 gals
Bees, bed bugs, boxelder bugs, fleas (outdoor), cecid flies, cockroaches, elm leaf beetles, ground beetles, moths, midges, crickets, flies (including such flies as stable, house, cluster and horseflies), hornets, killer bees, lice, mosquitoes, multicolored asian lady beetle, pillbugs, scorpions, saw-bugs, spiders, ticks (outdoor), yellowjackets, wasps	0.03% - 0.06%w/w	0.75-1.5 fl oz/gal 37.5-75 fl oz/50 gals 75-150 fl oz/100 gals
Termites (formosan & Eastern subterranean - localized control only)	0.06% w/w	1.5 fl oz/ gal 75 fl oz/50 gals 150 fl oz/100 gals
Carpenter ants	0.03% w/w	0.75 fl oz/gal 37.5 fl oz/50 gals 75 fl oz/100 gals

1 The recommended rate for maintenance treatments is 0.01% and for cleanout treatments is 0.03%. For control of severe infestations and longer residual control use the 0.06% rate. In addition, more frequent applications may be needed on concrete surfaces.

- 2 0.25 fl. oz = 7.5 milliliters  
0.75 fl. oz = 22.7 milliliters  
1.5 fl. oz = 45.4 milliliters

## GENERAL PEST CONTROL INDOORS

Treatment Sites: When used in accordance with label directions, Suspend SC Insecticide may be applied in and around residential and non-residential buildings and structures including residential kitchens and all types of food/feed handling establishments. Suspend SC Insecticide is intended to be mixed with water and applied indoors as a crack and crevice, spot or general surface treatment with hand pressurized or power operated sprayers or other suitable equipment for applying insecticides to localized areas. For residential buildings and structures apply only as a crack and crevice or spot treatment. This product can be applied to walls, floors, ceilings, in and around cabinets, between, behind and beneath equipment and appliances, around floor drains, window and door frames and around plumbing, sinks and other possible pest harborage sites. Permitted areas of use include, but are not limited to, the food/feed and non-food/non-feed areas of:

Aircraft (cargo only)	Food service establishments	Nursing homes
Apartment buildings	Granaries & grain mills	Offices
Bakeries	Hospitals	Railcars
Bottling facilities	Hotels	Restaurants
Breweries	Houses	Schools
Buses	Industrial buildings	Ships & vessels
Cafeterias	Installations	Trailers
Candy plants	Kitchens	Trucks
Canneries	Laboratories	Warehouses
Dairy product processing plants	Mausoleums	Wineries
Food manufacturing plants	Meat, poultry & egg processing & packaging plants	
Food processing plants	Mobile & motor homes	

Treatments may be applied at 21-day intervals or as necessary to maintain adequate control. Remove pets and cover aquariums and terrariums before spraying. Do not apply to classrooms when in use. Do not use in aircraft cabins.

Treat specific areas that insect inhabit or where they are frequently found. Use 0.25 to 1.5 fluid ounces of Suspend SC Insecticide in sufficient amount of water and apply to adequately cover 1000 square feet. Do not allow dripping or run-off to occur. Under conditions of severe pest infestation, when quick knockdown and/or longer residual control is desired, use 1.5 ounces of 0.06% Suspend SC Insecticide per 1,000 sq. ft.

**Premise Pest Control:** *Cockroaches, Ants, Crickets, Silverfish, Firebrats, Centipedes, Millipedes, Clothes Moths, Earwigs, Pillbugs, Sowbugs, Scorpions and Spiders:* Apply as a coarse, low pressure spot or crack and crevice spray to areas that these pests normally inhabit. Pay particular attention to dark corners of rooms and closets; floor drains; cracks and crevices in walls; along and behind baseboards; beneath and behind sinks, stoves, refrigerators, and cabinets; around plumbing and other utility installations; doors, windows and in attics and crawl spaces. Applications may be made to floor surfaces along baseboards and around air ducts. Let surfaces dry before allowing children or pets to contact surfaces.

**Pantry Pest Control:** To kill adult and accessible stages of insect pests such as *Lesser Grain Borers, Grain Weevils, Flour Beetles, Saw-toothed Grain Beetles, Cigarette Beetles, Merchant Grain Beetles, Angoumois Grain Moths, Mediterranean Flour Moths, Indian Meal Moths, Grain Mites and Spider Mites:* Make applications to cupboards, shelving and storage areas. Remove all foodstuffs, all utensils, shelf paper and other objects before spraying. Allow all treated surfaces to dry and cover shelves with clean paper prior to replacing foodstuffs, utensils or other items. Any foodstuff accidentally contaminated with any spray solution should be discarded.

**Crawling and Flying Insect Pests (such as flies, gnats, midges, mosquitoes, cockroaches, ants and silverfish):** Apply directly to walls, ceilings, window screens and other resting areas as a residual surface treatment. May be used inside residential buildings, as well as in and around carports, garages, storage sheds, unoccupied horse stables and pet kennels.

**Ants (including Pharaoh Ants):** Apply to trails, around doors and windows and other areas where ants inhabit or enter premises.

**Bedbugs and Lice:** Remove linens and wash before reuse. Apply to mattresses, especially tufts, folds, and edges until moist. Apply to the interior of the frame. Allow to dry before remaking bed.

**Carpet Beetles, Fleas and Ticks:** A total integrated pest management program is recommended. It is suggested that infested areas be thoroughly vacuumed prior to treatment. These areas include all carpet, upholstered furniture, baseboards and other indoor areas frequented by pets. The vacuum cleaner bag should be discarded in an outdoor trash container upon completion. Pet bedding should be cleaned or replaced. Apply Suspend SC Insecticide as a uniform coarse low-pressure spot or crack and crevice treatment to areas where these pests normally inhabit. Care should be taken to limit spot treatments to pet bedding and its surrounding area, infested floor coverings and pest habitats. Do not treat entire area of floor, carpet or floor coverings. Use a dust such as DeltaDust® in areas around electrical equipment. Do not apply as a space spray. Restrict applications to furniture where no prolonged contact by humans will occur. Do not allow people or pets to contact treated areas until spray has dried. This product can be used with an Insect Growth Regulator (IGR). Effective flea control includes controlling the source of flea infestations. This can require treatment of pets with a product registered by EPA for such use and treating outside areas frequented by pets. Suspend SC Insecticide can also be applied out-of-doors for flea control. Do NOT treat pets with this product.

**Pet Kennels:** For residual control of crawling and flying insects in and around these structures, apply as a general surface and/or crack and crevice spray. Optimal control results if facilities to be treated are clean and if the exterior perimeter is concurrently treated with Suspend SC Insecticide. Apply only to areas of facility where animals are not present. Allow spray to dry before allowing re-entry of animals into treated areas. Do not apply to animal feed or watering equipment.

**Food/Feed Handling Establishments:** For control of all labeled pests, applications are permitted in food/feed and non-food/non-feed areas of food/feed handling and processing establishments as a general surface, spot or crack and crevice treatment.

Food/Feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served. Included also are areas for receiving, storing, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food/feed. Serving areas where food/feed is exposed and the facility is in operation are also considered food/feed areas.

**Non-Food/Non-Feed Areas:** Examples of non-food/non-feed areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, mop closets and storage (after packaging, canning or bottling). All areas that insects inhabit or through which insects may enter should be treated.

**General Surface Applications:** Do not use this application method in the food/feed handling areas of the facility when the facility is in operation or foods/feeds are exposed. Do not apply directly to food/feed products. Cover or remove all food/feed processing and/or handling equipment during application. After application in food/feed processing plants, cafeterias and similar facilities, wash all equipment, benches, shelving and other surfaces which food/feed will contact. Clean food/feed handling or processing equipment and thoroughly rinse with clean, fresh water. After application in bakeries, clean all food contact surfaces thoroughly with appropriate equipment.

**Spot or Crack & Crevice Applications:** Except in federally inspected meat and poultry plants, spot or crack and crevice applications may be made while the facility is in operation provided exposed food is covered or removed from area being treated prior to application. Do not apply directly to food or food-handling surfaces.

In the home, do not allow spray to contact food/feed surfaces. If it does, clean surfaces with soap and water.

**GRAIN BINS AND WAREHOUSES:** Dilute 0.25-1.5 fl oz of Suspend SC Insecticide with water to make 1 gallon of solution. Apply finished spray to equipment, wall and floor surfaces of grain bins and warehouses at the rate of 1 gallon per 1000 sq. ft. prior to storing or handling grain. Do not allow run-off to occur. Application should be made only after equipment bins and buildings have been thoroughly cleaned. Empty bins or warehouses should be fumigated with an EPA registered fumigant according to the manufacturer's label recommendations when inaccessible areas exist. To help prevent infestation of these structures, apply Suspend SC Insecticide as a perimeter treatment where pests are active and may find entrance. It is recommended to treat a band of soil or lawn up to 6-10 feet wide around and adjacent to structures and treat the foundation to a height of 2 to 3 feet. Mix 0.25-1.5 fl oz of Suspend SC Insecticide with water to make 1 gallon of solution. Apply as a coarse spray at a rate of 1 gallon per 1000 sq. ft. Thoroughly and uniformly wet the barrier area but do not allow run-off to occur.

## GENERAL PEST CONTROL OUTDOORS

Use Suspend SC Insecticide to control outdoor pests by application as either a perimeter treatment,

residual spray or broadcast application to lawns, grounds and ornamental plantings.

**Treatment Sites:** When used in accordance with label directions, Suspend SC Insecticide may be applied as a perimeter treatment or as a residual spray around residential and non-residential settings. Begin applications when pests are noticed.

**Residual Spray:** Apply Suspend SC Insecticide as a coarse low-pressure spray at the rate of 0.75 to 1.5 fl oz to surfaces of buildings, porches, screens, window frames, eaves, patios, refuse dumps, garages and lawns.

**Perimeter Treatments:** To help prevent infestation of buildings, apply Suspend SC Insecticide where pests are active and may find entrance. It is recommended to treat a band of soil or lawn up to 6 - 10 feet wide around and adjacent to buildings and treat the building foundation to a height of 2 to 3 feet. Apply as a coarse spray at the rate of 0.25 to 1.5 fl oz of Suspend SC Insecticide in sufficient amount of water. (See Mixing Directions for Perimeter Application below.) Use an adequate volume to thoroughly and uniformly wet the barrier area but do not allow run-off to occur.

**Fleas and Ticks:** Apply Suspend SC Insecticide as a coarse wet spray to areas where fleas and ticks are most likely to be found, such as but not limited to, dog houses, kennels, runways, cracks and crevices in sidewalks, paths or patios, under plants, shrubs, bushes, trees, or other shaded, moist areas where pests may rest.

**Fire Ants:** Apply Suspend SC Insecticide as a drench. Dilute 1.5 fl oz per one gallon of water. Gently sprinkle 1 to 2 gallons of the diluted insecticide over the surface of each mound and surrounding area to a 2-foot diameter. For best results, apply in cool weather, 65 to 80°F or in early morning or late evening hours. Treat new mounds as they appear. Pressurized sprays may disturb the ants and cause migration, reducing product effectiveness.

**Wood-Infesting Insects:** For control of wood-infesting beetles on surfaces in homes and other structures, apply a 0.06% suspension of Suspend SC Insecticide. For treatment of small areas, apply by brushing or spraying the suspension evenly on wood surfaces. For large or overhead areas of homes, apartment buildings, etc., apply as a coarse spray to thoroughly cover the area. Cover all surfaces below the area being sprayed with plastic sheeting or other material which can be disposed of

by placing in trash if contamination from dripping occurs. Sprayed surfaces should be avoided until the treated area is completely dry.

For control of above-ground termites and wood-infesting beetles in localized areas, apply a 0.06% suspension or sufficient volume of foam of Suspend SC Insecticide to voids and galleries in damaged wood and in spaces between wooden structural members and between the sill plate and foundation where wood is vulnerable. Applications may be made to inaccessible areas by drilling and then injecting the suspension or foam with a suitable directional injector into the damaged wood or wall voids. All treatment holes drilled in construction elements in commonly occupied areas of structures must be securely plugged.

Termite carton nests in building voids may be injected with 0.06% suspension or foam using a pointed injection tool. Multiple injection points to varying depths may be necessary. It is desirable to physically remove carton nest material from building voids when such nests are found.

Application to attics, crawl spaces, unfinished basements, or man-made voids may be made with a coarse fan spray of 0.06% suspension to control exposed worker and winged reproductive forms of termites. For control of swarming or reproductive stages of termites, apply a 0.06% suspension to the swarm or area on which they collect.

The purpose of such applications of Suspend SC Insecticide is to kill workers or winged reproductive forms which may be present at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but are merely a supplement.

Do not apply solution until location of heat or air conditioning ducts, vents, water and sewer lines and electrical conduits are known and identified. When located, applications may be made around these structures. However, caution must be taken to avoid contamination of these structural elements and airways.

For control of above-ground termites inside trees, utility poles, fencing and decking materials and similar structural members, drill to find the interior infested cavity and inject 0.06% suspension or foam using an appropriate treatment tool with a splashback guard.

### MIXING DIRECTIONS FOR Suspend SC INSECTICIDE - OUTDOOR PERIMETER APPLICATION

#### I. Suspend SC INSECTICIDE DILUTION (0.25 fl oz/1000 sq. ft.) - (Based on one home being 1,000 sq. ft.)

Tank Size	15 Gallons		25 Gallons		50 Gallons		100 Gallons	
	Finished Spray: Gallons Per 1000 sq. ft.	Suspend SC Insecticide Per Tank (fl oz)	Homes Per Tank*	Suspend SC Insecticide Per Tank (fl oz)	Homes Per Tank*	Suspend SC Insecticide Per Tank (fl oz)	Homes Per Tank*	Suspend SC Insecticide Per Tank (fl oz)
1	3.75	15	6.25	25	12.50	50	25.00	100
3	1.25	5	2.08	8	4.17	16	8.33	33
5	0.75	3	1.25	5	2.50	10	5.00	20
10	0.37	1.5	0.63	2.5	1.25	5	2.50	10

\*Based on Suspend SC Insecticide applied at 0.25 ounces per 1,000 square feet treated per home or structure.

#### II. Suspend SC Insecticide dilution (1.00 fl oz/1000 sq. ft.)

Tank Size	15 Gallons		25 Gallons		50 Gallons		100 Gallons	
	Finished Spray: Gallons Per 1000 sq.ft.	Suspend SC Insecticide Per Tank (fl oz)	Homes Per Tank*	Suspend SC Insecticide Per Tank (fl oz)	Homes Per Tank*	Suspend SC Insecticide Per Tank (fl oz)	Homes Per Tank*	Suspend SC Insecticide Per Tank (fl oz)
1	15.0	15	25.0	25	50	50	100	100
3	5.0	5	8.4	8.3	16.70	16.7	33.3	33.3
5	3.0	3	5.0	5	10.0	10	20.0	20
10	1.5	1.5	2.5	2.5	5.0	5	10.0	10

\*Based on Suspend SC Insecticide applied at 1.00 ounce per 1,000 square feet treated per home or structure.

Termite carton nests in trees may be injected with 0.06% suspension or foam using a pointed injection tool. Multiple injection points to varying depths may be necessary. It is desirable to physically remove carton nest material from trees when such nests are found.

For protection of firewood from carpenter ants and termites, treat soil beneath firewood prior to stacking with a 0.06% suspension at 1 gallon per 8 square feet to prevent carpenter ant or termite infestation. DO NOT treat firewood with this product.

For control of carpenter ants in houses and other structures, use a 0.06% suspension and apply as a general surface, spot, crack and crevice or wall void application. Apply around doors and windows and other places where carpenter ants may enter the premises. Spray or foam into cracks and crevices through small drilled holes into voids where these ants or their nests are present. Use a sufficient amount of coarse spray or foam to cover the area. Do not exceed 1 gallon of dilute suspension per 1000 square feet of treated surface.

For control of carpenter ants inside trees, utility poles, fencing and decking materials and similar structural members, drill to find the interior infested cavity and inject 0.06% suspension or foam using an appropriate treatment tool with a splashback guard.

For control of tunneling carpenter ants in soil, apply a 0.06% suspension as a drench or inject the suspension or foam at intervals of 8 to 12 inches. Establish a uniform vertical barrier at the edge of walls, driveways or other hard surfaces where ants are tunneling beneath the surfaces.

For control of carpenter bees in structures, apply a 0.06% suspension or sufficient volume of foam directly into gallery entrance holes. Following treatment, the entrance holes may be left open 24 hours to be certain that returning adult bees are killed. When there is no activity, the hole may be sealed with a suitable sealant.

**Foam Application Use Directions:** Suspend SC Insecticide can be converted to a foam and the foam used to treat structural voids to control insect and arthropod pests in walls, under slabs or in other void areas. Mix end-use dilution of Suspend SC Insecticide with manufacturer's recommended volume of foaming agent. Use the recommended rate suspension converted to a foam to achieve the desired foam expansion ratios. Verify before treatment that the foaming agent is compatible with Suspend SC Insecticide.

**Turfgrass Applications:** Suspend SC Insecticide controls a broad spectrum of insects on turfgrasses and ornamental plants in outdoor areas such as athletic fields, recreational areas, residential landscapes, parks, and public and commercial sites such as around institutional commercial and industrial buildings.

Uniform coverage is essential. Begin applications when pests are evident. Apply Suspend SC Insecticide as a broadcast application using a conventional hydraulic boom-type sprayer or other suitable equipment for turfgrass pests according to the rates listed in the Application Rates for Turfgrass Sites table. Apply using a minimum of 2 gallons of water per 1000 sq. ft. Higher volumes (4 to 10 gallons of water per 1000 sq. ft.) may be needed for adequate coverage of dense turfgrass canopies and/or turfgrasses maintained under high mowing heights. Allow sufficient time for the application to dry before allowing pets or people to enter treatment area.

Target insects must be exposed to the insecticide to be controlled. For optimum control of surface-feeding insects, delay irrigation or mowing for 24 hours after application. For subsurface-feeding insects, irrigate with 0.25 to 0.5 inch of water immediately after application.

**Tank Mixing:** Suspend SC Insecticide has been found to be compatible with many commonly used pest control and fertilizer products. When tank-mixing Suspend SC Insecticide, observe all restrictions and precautions and limitations which appear on the label of each product. For any tank-mixture which has not been previously tested, mix the correct proportion of products and water in a quart jar to ensure proper physical compatibility of the mixture. Tank mixtures of Suspend SC Insecticide with silicone surfactants may improve the control of some insects.

**NOTE: In New York State, product may not be applied to turf within 100 feet of a coastal marsh or stream that drains directly into a coastal marsh. For control of turf insects in New York, a second application may be made after two weeks if target insect activity persists.**

## Application Rates For Turfgrass Sites:

Sites	Pests	Rates	Remarks
Athletic fields, recreational areas, lawns & grounds, residential areas, parks, public and commercial sites	Ants, armyworms, centipedes, chiggers, cockroaches, crickets, cutworms, fleas(adult), grasshoppers, ground beetles, Japanese beetles (adults), leafhoppers, mealybugs, pillbugs, scorpions, sod webworms, sowbugs, springtails, spiders, ticks	0.4 to 0.6 fl oz/1000 sq. ft or 17.5 to 26 fl oz/Acre or 0.06 to 0.08 lb ai/A	Apply when pests become evident. Apply as a broadcast spray in a minimum of 2 gallons of water per 1000 sq. ft. Delay irrigation or mowing for 24 hours after application for optimum control of surface-feeders such as cutworms and webworms. For subsurface - feeders, such as mole crickets, irrigate with 0.25 to 0.5 inch of water immediately after application.
	Annual bluegrass weevil (adult), billbugs (adult), black turfgrass ataenius (adult), chinch bugs, mites*, mole crickets.	0.6 to 0.9 fl oz/1000 sq. ft. or 26 to 39 fl oz/Acre or 0.08 to 0.13lb ai/A	The addition of a silicone surfactant may enhance mole cricket control.

### \*Suppression Only

### Landscape Ornamental Treatments:

Suspend SC Insecticide provides control of a wide range of insects on landscape ornamentals. Mix Suspend SC Insecticide with the required amount of water according to the Application Rates for Ornamentals table below. Thoroughly apply the finished spray to upper and lower leaf surfaces to ensure adequate coverage, but prior to dripping. Be sure to penetrate dense foliage to wet leaves, trunk and branches.

Plant safety testing has been conducted on many species of ornamental plants under a wide range of environmental and cultural conditions and no injury has been observed. If local use information is not available, it is advisable to pre-spray a few plants.

## Application Rates For Ornamentals:

Sites	Pests	Rates	Remarks
Landscape ornamentals including trees, shrubs, evergreens and foliage plants	Adelgids, ants, aphids, azalea lace bugs, bagworms, boxelder bugs, cankerworms, cardamom thrips, cockroaches, crickets, elm leaf beetles, elm spanworms, European pine sawflies, fall webworms, flea beetles, flies, fungus gnats, grasshoppers, greenstriped mapleworm, Gypsy moth (larvae), imported willow leaf beetles, Japanese beetles (adults)*, June beetles (adults), lace bugs, leaf-feeding caterpillars, leafhoppers, leaf rollers, loopers, leaf skeletonizers, mealybugs, midges, mimosa webworms, mites**, orangestriped oakworms, pillbugs, pine shoot beetles, pine tip moths, pinyon spindlegall midges, plant bugs, rehdumped caterpillar, red imported fire ants, sawfly larvae, sowbugs, spiders, springtails, spanworms, scale insects (crawlers), tent caterpillars, ticks, webworms, yellownecked caterpillar	4 to 8 oz/100 gallons of water	Apply when pests become evident. Thorough coverage is required for best results. Apply with sufficient water to ensure coverage of upper and lower leaf surfaces.

\*Will provide anti-feedant properties on treated ornamentals against Japanese beetles.

\*\*Suppression only.

### NOTES:

1. DO NOT apply Suspend SC Insecticide through any type of irrigation system.
2. DO NOT apply this product to edible crops.
3. DO NOT apply this product in hospital patient rooms while occupied or in any rooms occupied by the elderly or infirm.
4. DO NOT apply in classrooms when in use.
5. DO NOT apply to institutions (including libraries, sport facilities, etc.) in the immediate area when occupants are present.

## STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Mix as needed. Store in original container in a secured dry storage area. Avoid extremes in temperatures. Prevent cross-contamination with other pesticides and fertilizers. If the container is leaking and/or material is spilled, on floor or paved surfaces, absorb on sawdust or other commercially available absorbing material, sweep up and remove to chemical waste area for your disposal. Concentrate is stable at normal storage temperatures.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

## IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer (reg'd), the Bayer Cross (reg'd), Suspend®, and Backed by Bayer™ are trademarks of Bayer.



Produced for:

**Bayer Environmental Science**

A Division of Bayer CropScience LP

2 T. W. Alexander Drive

Research Triangle Park, NC 27709

Product of India